

CLAIM AMENDMENTS

---

1. *(Original)* A method for preparing cells for therapeutic use, comprising:  
a) differentiating human pluripotent stem (hPS) cells into a first cell population; and  
b) differentiating human pluripotent stem (hPS) cells into a second cell population;  
wherein the first cell population is MHC compatible with the second cell population, and  
whereupon administration of the first population to an individual renders the individual  
immunotolerant to the second cell population.
2. *(Original)* The method of claim 1, wherein the first cell population and the second cell population  
are differentiated from the same hPS cells or their progeny.
3. *(Original)* The method of claim 1, wherein the first cell population predominantly comprises  
mesoderm cells.
4. *(Original)* The method of claim 1, wherein the first cell population has characteristics of  
hematopoietic progenitor cells, blood leukocytes, leukocyte precursor cells, macrophage-like  
cells, dendritic cells, or mesenchymal stem cells.
5. *(Currently amended)* The method of claim 1, wherein the first cell population expresses one or  
more of the following markers: CD34, T-cell receptor, HLA Class II, CMRF-44, CMRF-56,  
DEC-205, S100, CD44, CD90, or CTLA-4.
6. *(Currently amended)* A method for preparing a first cell population that renders an individual to  
whom it is administered immunotolerant to a second cell population, comprising differentiating  
human pluripotent stem (hPS) cells into a mixed cell population, and enriching from the mixed  
population cells that express CD34, T-cell receptor, HLA Class II, CMRF-44, CMRF-56,  
DEC-205, S100, CD44, CD90, or CTLA-4.
7. *(Original)* The method of claim 1, wherein the second cell population comprises one of the  
following cell types or their lineage-restricted precursors: hepatocytes, neurons,  
oligodendrocytes, astrocytes, cardiomyocytes, or osteogenic cells.
8. *(Original)* A combination of pharmaceutical compounds, comprising in separate containers:  
a) a first cell population that has been differentiated from human pluripotent stem (hPS) cells  
into a phenotype that renders a subject to whom it is administered immunotolerant to a second  
cell population; and

b) the second cell population that is MHC compatible with the first cell population.

9. *(Original)* The pharmaceutical compounds of claim 8, wherein the first cell population and the second cell population are differentiated from the same hPS cell line.
10. *(Original)* The pharmaceutical compounds of claim 8, wherein the first cell population predominantly comprises mesoderm cells.
11. *(Original)* The pharmaceutical compounds of claim 8, wherein the first cell population has characteristics of hematopoietic progenitor cells, blood leukocytes, leukocyte precursor cells, macrophage-like cells, dendritic cells, or mesenchymal stem cells.
12. *(Currently amended)* The pharmaceutical compounds of claim 8, wherein the first cell population expresses one or more of the following markers: CD34, T-cell receptor, HLA Class II, CMRF-44, CMRF-56, DEC-205, S100, CD44, CD90, or CTLA-4.
13. *(Original)* The pharmaceutical compounds of claim 8, wherein the second cell population comprises one of the following cell types or their lineage-restricted precursors: hepatocytes, neurons, oligodendrocytes, astrocytes, cardiomyocytes, or osteogenic cells.
14. *(Original)* A method for reconstituting cellular function in an individual, comprising administering to the individual a first cell population and a second cell population, both differentiated from human pluripotent stem (hPS) cells, wherein the first cell population is MHC compatible with the second cell population,  
whereupon administration of the first cell population renders the individual immunotolerant to the second cell population; and  
whereupon administration of the second cell population reconstitutes the cellular function.
15. *(Currently amended)* The method of claim 14, wherein the phenotype of the first cell population expresses one or more of the following markers: CD34, T-cell receptor, HLA Class II, CMRF-44, CMRF-56, DEC-205, S100, CD44, CD90, or CTLA-4.
16. *(Original)* The method of claim 14, wherein the first cell population is administered to the circulation.
17. *(Original)* The method of claim 14, wherein the cellular function that is reconstituted in the individual is the function of hepatocytes, neurons, oligodendrocytes, astrocytes, cardiomyocytes, or osteogenic cells.

18. *(Original)* The method of claim 14, wherein administration of the second cell population occurs at least 2 weeks after administration of the first cell population.
19. *(Original)* The method of claim 14, wherein the first cell population and the second cell population are differentiated from the same hPS cells or their progeny.
20. *(Original)* A method of preparing an individual for therapy to reconstitute their cellular function, comprising administering to the individual a first cell population differentiated from human pluripotent stem (hPS) cells, thereby rendering the individual immunotolerant to a second cell population also differentiated from hPS cells that is MHC compatible with the first cell population, wherein the therapy comprises administering to the individual the second cell population, thereby reconstituting the cellular function in the individual.
-